

Abstract Submitted to the  
International Conference on Strongly Correlated Electron Systems  
University of Michigan, Ann Arbor  
August 6-10, 2001

### **Spin wave excitations in single crystalline $\text{U}_3\text{Pd}_{20}\text{Si}_6$**

Naofumi ASO<sup>1</sup>, Naoto METOKI<sup>2</sup>, Masahumi KOHGI<sup>3</sup>, Keith A. McEwen<sup>4</sup>, Yoshihiro KOIKE<sup>\*2</sup>, Yoshinori HAGA<sup>2</sup>, Naoyuki TATEIWA<sup>5</sup>, Noriaki KIMURA<sup>6</sup>, Haruyoshi AOKI<sup>6</sup>, Takemi KOMATSUBARA<sup>6</sup>, Yukio MORII<sup>2</sup>

<sup>1</sup> *NSL-ISSP, University of Tokyo, Tokai, Ibaraki 319-1106, Japan*

<sup>2</sup> *ASRC, Japan Atomic Energy Research Inst., Tokai, Ibaraki 319-1195, Japan*

<sup>3</sup> *Graduate School of Sci., Tokyo Metropolitan University, Hachioji, Tokyo 192-0397, Japan*

<sup>4</sup> *University College London, London WC1E 6BT, United Kingdom*

<sup>5</sup> *Research Center for Materials Science at Extreme Condition, Osaka University, Toyonaka, Osaka 560-8531, Japan*

<sup>6</sup> *Center for Low Temp. Sci., Tohoku University, Sendai 980-8545, Japan*

Inelastic neutron scattering were performed on single crystalline  $\text{U}_3\text{Pd}_{20}\text{Si}_6$  to investigate the nature of the 5f electrons of uranium compounds. Antiferromagnetic spin wave excitations due to the uranium 5f electrons at the 8c site were observed below the Néel temperature  $T_N = 19$  K, exhibiting the localized nature of the 5f electrons. Its line width was found to show the energy resolution limited behaviour except for the the zone center. The anomolous line width around the zone center will be discussed in connection with the magnetic ordering at the 4a site below  $T = 2$  K.

---

<sup>\*</sup>Japan Science and Technology Corporation, Kawaguchi, Saitama 332-0012, Japan